



**MASSACHUSETTS DEPARTMENT OF PUBLIC HEALTH**  
**WEEKLY INFLUENZA UPDATE**  
**December 19, 2013**

All data in this report are preliminary and subject to change as more information is received.


**Sentinel Provider Surveillance: Influenza-like illness activity**

Week 50 Activity<sup>1</sup> (representing geographic distribution): Regional

Week 50 ILI Activity<sup>2</sup> (representing intensity of ILI activity): 2 (Minimal)

Provider offices across the US report the amount of influenza-like illness (ILI) they see in their patients each week during regular flu season. These outpatient providers' offices, which include doctors' offices, school health services, and community health centers, are called 'sentinel sites.' Here we present Massachusetts sentinel site data. Please note that the data represent not only confirmed influenza cases, but also those just with ILI, which may be caused by other viruses. ILI is defined as fever above 100.0<sup>1</sup> in addition to either cough or sore throat. ILI is a marker of influenza and is used throughout the regular influenza season to monitor influenza since most people are not tested for influenza. Figure 1 shows a very modest increase in ILI activity since the beginning of the season, while there is elevated activity in other regions of the United States. For more information, see CDC's influenza surveillance website at [www.cdc.gov/flu/weekly/fluactivitysurv.htm](http://www.cdc.gov/flu/weekly/fluactivitysurv.htm).

**Figure 1: Percentage of ILI visits reported by sentinel provider sites**



\*Influenza-like illness (ILI, defined by fever >100F and cough and/or sore throat), as reported by Massachusetts sentinel surveillance sites.

<sup>1</sup> CDC activity indicator – indicates how widespread influenza activity level is in the state.

<sup>2</sup> CDC ILI activity indicator – more quantitative indicator of the level of ILI activity across the state.

Table 1 below shows a geographical distribution of reported ILI in Massachusetts. Table 1 shows that sentinel sites in three regions of the state are experiencing elevated ILI activity.

**Table 1: Percent ILI reported weekly by Massachusetts sentinel sites**


	Regional Baseline % ILI*	2013-2014			2012-2013		
		% ILI	Report. Sites	Total enroll.	% ILI	Report. Sites	Total enroll.
<b>Boston</b>	0.95	0.83	6	6	2.06	6	5
<b>Central</b>	1.35	1.04	9	12	1.29	11	11
<b>Inner Metro Boston</b>	0.88	1.09	8	13	2.04	11	13
<b>Northeast</b>	0.92	1.49	9	11	1.69	11	11
<b>Outer Metro Boston</b>	1.32	1.42	4	5	4.54	4	4
<b>Southeast</b>	1.43	0.28	3	6	1.48	4	3
<b>West</b>	1.36	0.52	7	7	1.50	5	7

\*Regional baseline % ILI is calculated weekly using reporting providers' baseline % ILI estimates.

**Influenza-Associated Hospitalizations**

In 2010, MDPH began to request voluntary reporting of all laboratory-confirmed influenza hospitalizations from hospitals in Massachusetts. As many as 50 acute care hospitals from across the state report these data to MDPH on a weekly basis during flu season. The graph below shows the number of laboratory-confirmed hospitalizations per 1,000 licensed beds represented by reporting hospitals for the current season and two previous seasons.


**Figure 2: Massachusetts laboratory-confirmed influenza hospitalizations**



**Laboratory testing for influenza**

Laboratories in Massachusetts report all positive influenza laboratory tests to MDPH, including viral culture, polymerase chain reaction (PCR) and rapid influenza diagnostic tests. Because the majority of cases are not tested, the number of 'confirmed' cases does not reflect the overall incidence of influenza; however, this information is essential to track the types of influenza circulating in Massachusetts and can be a useful indicator of the presence and distribution of influenza in the state. Figure 3 illustrates the number of laboratory confirmed cases in Massachusetts by week, shown along with Massachusetts ILI. Table 2 reflects the number of laboratory-confirmed influenza cases by region and influenza type.

**Figure 3: Laboratory-confirmed Influenza Cases and Influenza-like Illness  
Massachusetts, September 29, 2013 – December 18, 2013**



\*Influenza cases confirmed via viral culture, PCR or rapid test by specimen collection date.

\*\*Influenza-like illness (ILI, defined as fever>100F and cough and/or sore throat), as reported by Massachusetts sentinel surveillance sites by CDC week date.


**Table 2: Laboratory-confirmed Influenza by Region – 2013-2014 and 2012-2013 Influenza Seasons**

Region	2013-2014						2012-2013					
	A		B		Untyped		A		B		Untyped	
	Week	YTD	Week	YTD	Week	YTD	Week	YTD	Week	YTD	Week	YTD
Boston	8	34	0	11	0	0	141	265	3	9	1	2
Central	5	36	1	65	1	5	296	516	23	57	12	15
Inner Metro Boston	7	58	1	8	0	2	349	551	12	26	26	31
Northeast	36	130	11	73	6	16	392	568	9	50	73	95
Outer Metro Boston	12	34	1	6	0	2	231	362	25	52	28	38
Southeast	6	50	0	46	0	6	239	402	25	50	39	56
Unknown	3	6	0	6	0	0	46	94	11	24	0	1
West	1	11	0	1	0	0	56	105	18	26	1	1
<b>Total</b>	<b>78</b>	<b>359</b>	<b>14</b>	<b>216</b>	<b>7</b>	<b>31</b>	<b>1,750</b>	<b>2,863</b>	<b>126</b>	<b>294</b>	<b>180</b>	<b>239</b>

**Testing at the Hinton State Laboratory Institute**

MDPH’s Bureau of Laboratory Sciences (MDPH-BLS) performs confirmatory testing, typing and subtyping of influenza using PCR and viral culture primarily for samples submitted by ~60 outpatient healthcare providers (ILINet) as well as for early influenza positives and unusual cases from clinical hospital diagnostic laboratories across Massachusetts. For the 2013-2014 season, Figure 4 and Table 3 summarize virologic surveillance testing conducted by MDPH-BLS beginning MMWR week 40 (week ending October 5, 2013). MDPH-BLS performs influenza surveillance testing year round. One additional case of A (2009 H1N1) was identified in week 50, totaling five cases of A (2009 H1N1) two cases of A (H3N2) and one case of Flu B confirmed for the 2013-2014 season to date.

**Figure 4: Influenza positive tests reported to CDC by MDPH-BLS, September 29, 2013 – December 14, 2013**



**Table 3: Weekly Summary of MDPH-BLS Influenza Surveillance Test Results**

<b>2013-2014 Season: Influenza Surveillance</b>									
MA Department of Public Health's Bureau of Laboratory Sciences (MDPH-BLS)									
MMWR Week: (Specimen Collected)	2009 H1N1	seasonal A/H3N2	B	H3N2v	A unsub	No. Flu Pos (%)	Unsat	Total Tested	Total Rec'd
47 (11/17- 11/23/2013)	1	0	1	0	0	<b>2(33%)</b>	2	<b>6</b>	<b>8</b>
48 (11/24- 11/30/2013)	1	0	0	0	0	<b>1(33%)</b>	2	<b>3</b>	<b>5</b>
49 (12/01- 12/07/2013)	1	0	0	0	0	<b>1(50%)</b>	5	<b>2</b>	<b>7</b>
50 (12/08- 12/14/2013)	1	0	0	0	0	<b>1(10%)</b>	2	<b>10</b>	<b>12</b>
<b>Prior 4 wk Total</b>	<b>4</b>	<b>0</b>	<b>1</b>	<b>0</b>	<b>0</b>	<b>5(33%)</b>	<b>11</b>	<b>21</b>	<b>32</b>
<b>Cumulative Season total</b>	<b>5</b>	<b>2</b>	<b>1</b>	<b>0</b>	<b>0</b>	<b>8(16%)</b>	<b>15</b>	<b>49</b>	<b>64</b>

All data are subject to change. The 2013-2014 influenza season began MMWR 40 (09/29- 10/05/2013).

At the start of the 2013-2014 season, the first 10 isolates and thereafter 5 representative isolates every two weeks will be sent by MDPH-BLS to a CDC contract lab performing National Routine Surveillance to include antigenic characterization by hemagglutination inhibition (HI), genetic analysis (sequencing) and sensitivity to FDA-approved drugs for identification of resistance. As samples are received, MDPH-BLS will screen additional samples every two weeks to detect point mutations within the neuraminidase gene of H3N2 viruses (E119, R292, and N294) and influenza A (2009) viruses (H275 and I223) to assess resistance trends. This information will be reported locally and captured nationally in FluView.

There were three 2009 A(H1N1) isolates from MA during the 2009-2010 season with a mutation conferring oseltamivir-resistance (H275Y) and none during the 2010-2011, 2011-2012 and 2012-13 seasons. MDPH-BLS has not identified any isolates with antiviral resistance-associated mutations for the 2013-2014 season to date.

**Table 4: MDPH-BLS Influenza Antiviral Resistance Screening: 2013-2014 Season**

<b>Virus Collection Period: September 29, 2013- ongoing</b>				
	<b>Oseltamivir</b>		<b>Zanamivir</b>	
	<b>Samples Tested</b>	<b>Resistant Viruses, Number (%)</b>	<b>Samples Tested</b>	<b>Resistant Viruses, Number (%)</b>
<b>Influenza A (H3N2)<sup>i</sup></b>	2	0 (0)	2	0 (0)
<b>Influenza A (H1N1)pdm09<sup>ii</sup></b>	2	0 (0)	0	0 (0)

<sup>i</sup> Samples tested by pyrosequencing at position E119, R292, and N294 within the neuraminidase (NA) gene.

<sup>ii</sup> Samples tested by pyrosequencing at position H275 and I223 within the NA gene.

Additional information on national antiviral resistance testing can be found at <http://www.cdc.gov/flu/weekly/>. Recommendations for antiviral treatment and chemoprophylaxis of influenza virus infection can be found at <http://www.cdc.gov/flu/professionals/antivirals/index.htm>.